

- nation antiretroviral therapy including HIV protease inhibitors. *Aids*. 1998;12:613-8.
143. Martin DF, Sierra-Madero J, Walmsley S, Woltz RA, Macey K, Georgiou P, et al. A controlled trial of valganciclovir as induction therapy for cytomegalovirus retinitis. *N Engl J Med*. 2002;346:1119-26.
 144. Zegans ME, Walton RC, Holland GN, O'Donnell JJ, Jacobson MA, Margolis TP. Transient vitreous inflammatory reactions associated with combination antiretroviral therapy in patients with AIDS and cytomegalovirus retinitis. *Am J Ophthalmol*. 1998;125:292-300.
 145. Karavellas MP, Plummer DJ, Macdonald JC, Torriani FJ, Shufelt CL, Azen SP, et al. Incidence of immune recovery vitritis in cytomegalovirus retinitis patients following institution of successful highly active antiretroviral therapy. *J Infect Dis*. 1999;179:697-700.
 146. Spector SA, McKinley GF, Lalezari JP, Samo T, Andruczk R, Follansbee S, et al. Oral ganciclovir for the prevention of cytomegalovirus disease in persons with AIDS. Roche Cooperative Oral Ganciclovir Study Group. *N Engl J Med*. 1996;334:1491-7.
 147. Brosgart CL, Louis TA, Hillman DW, Craig CP, Alston B, Fisher E, et al. A randomized, placebo-controlled trial of the safety and efficacy of oral ganciclovir for prophylaxis of cytomegalovirus disease in HIV-infected individuals. Terry Bein Community Programs for Clinical Research on AIDS. *Aids*. 1998;12:269-77.
 148. Wohl DA, Zeng D, Stewart P, et al. Cytomegalovirus viremia, mortality, and end-organ disease among patients with AIDS receiving potent antiretroviral therapies. *J Acquir Immune Defic Syndrom*. 2005;38:538-44.
 149. Wohl D, Kendall M, Andersen J, Crumpacker C, Spector S, Feinberg J, et al. Randomized, placebo controlled trial of Valganciclovir to prevent CMV end-organ disease among HIV-infected subjects with detectable plasma CMV DNA-PCR: ACTG 5030 [abstract 150]. 14th Conference on Retroviruses and Opportunistic Infections. Boston (USA); 2006.
 150. Paltiel AD, Goldie SJ, Losina E, Weinstein MC, Seage GR 3rd, Kimmel AD, et al. Preevaluation of clinical trial data: the case of preemptive cytomegalovirus therapy in patients with human immunodeficiency virus. *Clin Infect Dis*. 2001;32: 783-93.
 151. Jacobson MA. Treatment of cytomegalovirus retinitis in patients with the acquired immunodeficiency syndrome. *N Engl J Med*. 1997;337:105-14.
 152. Drew WL, Ives D, Lalezari JP, Crumpacker C, Follansbee SE, Spector SA, et al. Oral ganciclovir as maintenance treatment for cytomegalovirus retinitis in patients with AIDS. Syntex Cooperative Oral Ganciclovir Study Group. *N Engl J Med*. 1995;333:615-20.
 153. Curran M, Noble S. Valganciclovir. *Drugs*. 2001;61:1145-50; discussion 1151-2.
 154. Musch DC, Martin DF, Gordon JF, Davis MD, Kuppermann BD. Treatment of cytomegalovirus retinitis with a sustained-release ganciclovir implant. The Ganciclovir Implant Study Group. *N Engl J Med*. 1997;337: 83-90.
 155. Martin DF, Kuppermann BD, Woltz RA, Palestine AG, Li H, Robinson CA. Oral ganciclovir for patients with cytomegalovirus retinitis treated with a ganciclovir implant. Roche Ganciclovir Study Group. *N Engl J Med*. 1999; 340:1063-70.
 156. Perry CM, Balfour JA. Fomivirsen. *Drugs*. 1999;57:375-80.
 157. Whitley RJ, Jacobson MA, Friedberg DN, Holland GN, Jabs DA, Dieterich DT, et al. Guidelines for the treatment of cytomegalovirus diseases in patients with AIDS in the era of potent antiretroviral therapy: recommendations of an international panel. International AIDS Society-USA. *Arch Intern Med*. 1998;158:957-69.
 158. Tural C, Romeu J, Sirena G, Andreu D, Conejero M, Ruiz S, et al. Long-lasting remission of cytomegalovirus retinitis without maintenance therapy in human immunodeficiency virus-infected patients. *J Infect Dis*. 1998;177: 1080-3.
 159. Vrabec TR, Baldassano VF, Whitchup SM. Discontinuation of maintenance therapy in patients with quiescent cytomegalovirus retinitis and elevated CD4+ counts. *Ophthalmology*. 1998;105:1259-64.
 160. Whitchup SM, Fortin E, Lindblad AS, Griffiths P, Metcalf JA, Robinson MR, et al. Discontinuation of anticytomegalovirus therapy in patients with HIV infection and cytomegalovirus retinitis. *JAMA*. 1999;282:1633-7.
 161. Macdonald JC, Torriani FJ, Morse LS, Karavellas MP, Reed JB, Freeman WR. Lack of reactivation of cytomegalovirus (CMV) retinitis after stopping CMV maintenance therapy in AIDS patients with sustained elevations in CD4 T cells in response to highly active antiretroviral therapy. *J Infect Dis*. 1998;177:1182-7.
 162. Jouan M, Saves M, Tubiana R, Carcelain G, Cassoux N, Aubron-Olivier C, et al. Discontinuation of maintenance therapy for cytomegalovirus retinitis in HIV-infected patients receiving highly active antiretroviral therapy. *Aids*. 2001;15:23-31.
 163. Berenguer J, González J, Pulido F, Padilla B, Casado JL, Rubio R, et al. Discontinuation of secondary prophylaxis in patients with cytomegalovirus retinitis who have responded to highly active antiretroviral therapy. *Clin Infect Dis*. 2002;34:394-7.
 164. Centers for Disease Control and Prevention, Workowski KA, Berman SM. Sexually transmitted diseases treatment guidelines, 2006. *MMWR Recomm Rep*. 2006;55:1-94.
 165. Nagot N, Ouédraogo A, Foulongne V, Konaté I, Weiss HA, Vergne L, et al. Reduction of HIV-1 RNA levels with therapy to suppress herpes simplex virus. *N Engl J Med*. 2007;356:790-9.
 166. Marin M, Güris D, Chaves SS, Schmid S, Seward JF; Advisory Committee on Immunization Practices, Centers for Disease Control and Prevention (CDC). Prevention of varicella: recommendations of the Advisory Committee on Immunization Practices (ACIP). *MMWR Recomm Rep*. 2007;56:1-40.
 167. González-García J, Guerra L, Amela C, del Amo J, Bruguera M, Castilla J, et al. Coinfección por VIH y virus de las hepatitis A, B y C en pacientes adultos. Revisión y recomendaciones de GESIDA/PNS. En: Grupo de trabajo para la elaboración de recomendaciones sobre las hepatitis virales en pacientes infectados por el VIH; 2002. Disponible en: www.gesidaseimc.com
 168. Weissman S, Feucht C, Moore BA. Response to hepatitis A vaccine in HIV-positive patients. *J Viral Hepat*. 2006;13:81-6.
 169. Launay O, Grabar S, Gordien E, Desaint C, Dimet J, Jegou D, et al. Randomized Trial Comparing a 3-Dose Regimen to a Standard 2-Dose Regimen of Hepatitis A Vaccine in HIV-infected Adult Patients with CD4+ T Cells between 200 and 500 per mm³ (HEPAVAC Study). *CROI 2007*. Los Angeles. February 25-27, 2007. Poster 885.
 170. Vento S, Garofano T, Renzini C, Cainelli F, Casali F, Ghironzi G, et al. Fulminant hepatitis associated with hepatitis A virus superinfection in patients with chronic hepatitis C. *N Engl J Med*. 1998;338:286-90.
 171. Fonseca MO, Pang LW, de Paula Cavalheiro N, Barone AA, Heloisa Lopes M. Randomized trial of recombinant hepatitis B vaccine in HIV-infected adult patients comparing a standard dose to a double dose. *Vaccine*. 2005; 23:2902-8.
 172. Keet IP, van Doornum G, Safary A, Coutinho RA. Insufficient response to hepatitis B vaccination in HIV-positive homosexual men. *AIDS*. 1992;6: 509-10.
 173. Rey D, Krantz V, Partisani M, Schmitt MP, Meyer P, Libbrecht E, et al. Increasing the number of hepatitis B vaccine injections augments anti-HBs response rate in HIV-infected patients. Effects on HIV-1 viral load. *Vaccine*. 2000;18:1161-5.
 174. Cooper CL, Davis HL, Angel JB, Morris ML, Elfer SM, Seguin I, et al. CPG 7909 adjuvant improves hepatitis B virus vaccine seroprotection in antiretroviral-treated HIV-infected adults. *AIDS*. 2005;19:1473-9.
 175. Cooper CL, Angel JB, Seguin I, Davis HL, Cameron DW. CPG 7909 adjuvant plus hepatitis B virus vaccination in HIV-infected adults achieves long-term seroprotection for up to 5 years. *Clin Infect Dis*. 2008;46:1310-4.
 176. Mast EE, Weinbaum CM, Fiore AE, Alter MJ, Bell BP, Finelli L, et al. A comprehensive immunization strategy to eliminate transmission of hepatitis B virus infection in the United States: recommendations of the Advisory Committee on Immunization Practices (ACIP) Part II: immunization of adults. *MMWR Recomm Rep*. 2006;55:1-33.
 177. Neuzil KM, Coffey CS, Mitchel EF, Jr., Griffin MR. Cardiopulmonary hospitalizations during influenza season in adults and adolescents with advanced HIV infection. *J Acquir Immune Defic Syndr*. 2003;34:304-7.
 178. Bridges CB, Fukuda K, Uyeki TM, Cox NJ, Singleton JA. Prevention and control of influenza. Recommendations of the Advisory Committee on Immunization Practices (ACIP). *MMWR Recomm Rep*. 2002;51:1-31.
 179. Berenguer J, Miralles P, Arrizabalaga J, Ribera E, Dronda F, Baraia-Etxaburu J, et al. Clinical course and prognostic factors of progressive multifocal leukoencephalopathy in patients treated with highly active antiretroviral therapy. *Clin Infect Dis*. 2003;36:1047-52.
 180. Selwyn PA, Hartel D, Lewis VA, Schoenbaum EE, Vermund SH, Klein RS, et al. A prospective study of the risk of tuberculosis among intravenous drug users with human immunodeficiency virus infection. *N Engl J Med*. 1989;320:545-50.
 181. Daley CL, Small PM, Schecter GF, Schoolnik GK, McAdam RA, Jacobs WR, Jr., et al. An outbreak of tuberculosis with accelerated progression among persons infected with the human immunodeficiency virus. An analysis using restriction-fragment-length polymorphisms. *N Engl J Med*. 1992;326: 231-5.
 182. Antiretroviral Therapy in Low-Income Countries Collaboration of the International epidemiological Databases to Evaluate AIDS (IeDEA); ART Cohort Collaboration. Tuberculosis after initiation of antiretroviral therapy in low-income and high-income countries. *Clin Infect Dis*. 2007;45:1518-21.
 183. Moreno S, Jarrin I, Iribarren JA, Viciano P, Parra J, Gomez-Sirvent JL, et al. Impact of Highly Active Antiretroviral Therapy (HAART) on the Incidence of Tuberculosis among HIV-Infected Patients. Presentado en: 14th Conference on Retroviruses and Opportunistic Infections. Los Angeles, 2007. Resumen 847.
 184. Muga R, Ferreros I, Langohr K, de Olalla PG, Del Romero J, Quintana M, et al. Changes in the incidence of tuberculosis in a cohort of HIV-seroconverters before and after the introduction of HAART. *AIDS*. 2007;21: 2521-7.

185. Luetkemeyer AF, Charlebois ED, Flores LL, Bangsberg DR, Deeks SG, Martin JN, et al. Comparison of an interferon- γ release assay with tuberculin skin testing in HIV-infected individuals. *Am J Respir Crit Care Med*. 2007;175:737-42.
186. Jones S, de Gijssel D, Wallach FR, Gurtman AC, Shi Q, Sacks H. Utility of QuantiFERON-TB Gold in-tube testing for latent TB infection in HIV-infected individuals. *Int J Tuberc Lung Dis*. 2007;11:1190-5.
187. Rangaka MX, Wilkinson DA, Seldon R, Van Cutsem G, Meintjes GA, Morrison C, et al. Effect of HIV-1 infection on T-cell-based and skin test detection of tuberculosis infection. *Am J Respir Crit Care Med*. 2007;175:514-20.
188. Lawn SD, Bangani N, Vogt M, Gail-Bekker M, Badri M, Ntobongwana M, et al. Utility of interferon- γ ELISPOT assay responses in highly tuberculosis-exposed patients with advanced HIV infection in South Africa. *BMC Infect Dis*. 2007;7:99-108.
189. Domínguez J, Ruiz-Manzano J, De Souza-Galvão M, Latorre I, Milà C, Blanco S, et al. Comparison of two commercially available gamma interferon blood tests for immunodiagnosis of tuberculosis. *Clin Vaccine Immunol*. 2008;15:168-71.
190. Pai M, Zwerling A, Menzies D. Systematic Review: T-Cell-Based Assays for the Diagnosis of Latent Tuberculosis Infection: An Update. *Ann Intern Med*. 2008 Jun 30. [Epub ahead of print].
191. Centers for Disease Control and Prevention. Anergy skin testing and tuberculosis preventive therapy for HIV-infected persons: revised recommendations. Centers for Disease Control and Prevention. *Morb Mortal Wkly Rep*. 1997;46:1-10.
192. Whalen CC, Johnson JL, Okwera A, Hom DL, Huebner R, Mugenyi P, et al. A trial of three regimens to prevent tuberculosis in Ugandan adults infected with the human immunodeficiency virus. Uganda-Case Western Reserve University Research Collaboration. *N Engl J Med*. 1997;337:801-8.
193. Gordin FM, Matts JP, Miller C, Brown LS, Hafner R, John SL, et al. A controlled trial of isoniazid in persons with anergy and human immunodeficiency virus infection who are at high risk for tuberculosis. Terry Beirn Community Programs for Clinical Research on AIDS. *N Engl J Med*. 1997;337:315-20.
194. Rivero A, López-Cortés L, Castillo R, Lozano F, García MA, Díez F, et al. Ensayo clínico aleatorizado de tres pautas de quimioprofilaxis para prevenir la tuberculosis en pacientes infectados por el VIH con anergia cutánea. *Enferm Infecc Microbiol Clin*. 2003;21:287-92.
195. Woldehanna S, Volmink J. Treatment of latent tuberculosis infection in HIV infected persons. *Cochrane Database Syst Rev*. 2004;1:CD000171.
196. Elzi L, Schelegel M, Weber R, Hirschel B, Cavassini M, Schmid P, et al. Reducing tuberculosis incidence by tuberculin skin testing, preventive treatment, and antiretroviral therapy in an area of low tuberculosis transmission. *Clin Infect Dis*. 2007;44:94-102.
197. Selwyn PA, Skellern BM, Alcabes P, Friedland GH, Klein RS, Schoenbaum EE. High risk of active tuberculosis in HIV-infected drug users with cutaneous anergy. *JAMA*. 1992;268:504-9.
198. Moreno S, Baraia-Etxaburu J, Bouza E, Parras F, Pérez-Tascón M, Miralles P, et al. Risk for developing tuberculosis among anergic patients infected with HIV. *Ann Intern Med*. 1993;119:194-8.
199. Guelar A, Gatell JM, Verdejo J, Podzamczar D, Lozano L, Aznar E, et al. A prospective study of the risk of tuberculosis among HIV-infected patients. *Aids*. 1993;7:1345-9.
200. Pape JW, Jean SS, Ho JL, Hafner A, Johnson WD, Jr. Effect of isoniazid prophylaxis on incidence of active tuberculosis and progression of HIV infection. *Lancet*. 1993;342:268-72.
201. Moreno S, Miralles P, Díaz MD, Baraia J, Padilla B, Berenguer J, et al. Isoniazid preventive therapy in human immunodeficiency virus-infected persons. Long-term effect on development of tuberculosis and survival. *Arch Intern Med*. 1997;157:1729-34.
202. Hawken MP, Meme HK, Elliott LC, Chakaya JM, Morris JS, Githui WA, et al. Isoniazid preventive therapy for tuberculosis in HIV-1-infected adults: results of a randomized controlled trial. *Aids*. 1997;11:875-82.
203. Mwinga A, Hosp M, Godfrey-Faussett P, Quigley M, Mwaba P, Mugala BN, et al. Twice weekly tuberculosis preventive therapy in HIV infection in Zambia. *Aids*. 1998;12:2447-57.
204. Rivero A, López Cortés L, Castillo R, Verdejo J, García MA, Martínez Marcos FJ, et al. Ensayo clínico aleatorizado para evaluar tres pautas cortas de tratamiento de la infección latente tuberculosa en pacientes infectados por VIH. *Enferm Infecc Microbiol Clin*. 2007;25:305-10.
205. Geijo MP, Herranz CR, Vaño D, García AJ, García M, Dimas JF. Pauta corta de isoniazida y rifampicina comparada con isoniazida para la infección latente de tuberculosis. Ensayo clínico aleatorizado. *Enferm Infecc Microbiol Clin*. 2007;25:300-4.
206. Gordin F, Chaisson RE, Matts JP, Miller C, de Lourdes García M, Hafner R, et al. Rifampin and pyrazinamide vs isoniazid for prevention of tuberculosis in HIV-infected persons: an international randomized trial. Terry Beirn Community Programs for Clinical Research on AIDS, the Adult AIDS Clinical Trials Group, the Pan American Health Organization, and the Centers for Disease Control and Prevention Study Group. *JAMA*. 2000;283:1445-50.
207. Halsey NA, Coberly JS, Desormeaux J, Losikoff P, Atkinson J, Moulton LH, et al. Randomised trial of isoniazid versus rifampicin and pyrazinamide for prevention of tuberculosis in HIV-1 infection. *Lancet*. 1998;351:786-92.
208. Ena J, Valls V. Short-course therapy with rifampin plus isoniazid, compared with standard therapy with isoniazid, for latent tuberculosis infection: a meta-analysis. *Clin Infect Dis*. 2005;40:670-6.
209. Martínez Alfaro E, Cuadra F, Solera J, Maciá MA, Geijo P, Sánchez Martínez PA y miembros del grupo GECMEI. Evaluación de dos pautas de quimioprofilaxis tuberculosa en pacientes infectados por el virus de la inmunodeficiencia humana. *Med Clin (Barc)*. 2000;115:161-5.
210. American Thoracic Society y Centers for Disease Control and Prevention. Targeted tuberculin testing and treatment of latent tuberculosis infection. *Am J Respir Crit Care Med*. 2000;161:S221-47.
211. Centers for Disease Control and Prevention. Update: adverse event data and revised American Thoracic Society/CDC recommendations against the use of rifampin and pyrazinamide for treatment of latent tuberculosis infection – United States, 2003. *MMWR Morb Mortal Wkly Rep*. 2003;52:735-9.
212. Gordin FM, Cohn DL, Matts JP, Chaisson RE, O'Brien RJ. Hepatotoxicity of rifampin and pyrazinamide in the treatment of latent tuberculosis infection in HIV-infected persons: is it different than in HIV-uninfected persons? *Clin Infect Dis*. 2004;39:561-5.
213. Ijaz K, Jereb JA, Lambert LA, Bower WA, Spradling PR, McElroy PD, et al. Severe or fatal liver injury in 50 patients in the United States taking rifampin and pyrazinamide for latent tuberculosis infection. *Clin Infect Dis*. 2006;42:346-55.
214. Burman WJ. Short-course regimens for latent tuberculosis: what is ready for prime time? *Enferm Infecc Microbiol Clin*. 2007;25:297-9.
215. Moreno S, Podzamczar D, Blazquez R, Iribarren JA, Ferrer E, Reparaz J, et al. Treatment of tuberculosis in HIV-infected patients: safety and antiretroviral efficacy of the concomitant use of ritonavir and rifampin. *Aids*. 2001;15:1185-7.
216. Oliva J, Moreno S, Sanz J, Ribera E, Molina JA, Rubio R, et al. Co-administration of rifampin and nevirapine in HIV-infected patients with tuberculosis. *Aids*. 2003;17:637-8.
217. Patel A, Patel K, Patel J, Shah N, Patel B, Rani S. Safety and antiretroviral effectiveness of concomitant use of rifampicin and efavirenz for antiretroviral-naïve patients in India who are coinfecting with tuberculosis and HIV-1. *J Acquir Immune Defic Syndr*. 2004;37:1166-9.
218. Pedral-Sampaio D, Alves C, Netto E, Brites C, Oliveira AS, Badaro R. Efficacy and safety of efavirenz in HIV patients on rifampin for tuberculosis. *Brazil J Infect Dis*. 2004;8:211-6.
219. Manosuthi W, Kiertiburanakul S, Ruxrungtham K, Sungkanuparph S, Vibhagool A, Rattanasiri S, et al. Efavirenz 600 mg/day versus efavirenz 800 mg/day in HIV-infected patients with tuberculosis receiving rifampicin: 48 week results. *AIDS*. 2006;20:131-2.
220. Centers for Disease Control and Prevention. Managing Drug Interactions in the Treatment of HIV-Related Tuberculosis [online]. 2007. Disponible en: <http://www.cdc.gov/tb/TB/HIV/Drugs/default.htm>
221. Weltman AC, Rose DN. The safety of Bacille Calmette-Guérin vaccination in HIV infection and AIDS. *Aids*. 1993;7:149-57.
222. Havlir DV, Dube MP, Sattler FR, Forthal DN, Kemper CA, Dunne MW, et al. Prophylaxis against disseminated *Mycobacterium avium* complex with weekly azithromycin, daily rifabutin, or both. *N Engl J Med*. 1996;335:392-8.
223. Pierce M, Crampton S, Henry D, Heifets L, LaMarca A, Montecalvo M, et al. A randomized trial of clarithromycin as prophylaxis against disseminated *Mycobacterium avium* complex infection in patients with advanced acquired immunodeficiency syndrome. *N Engl J Med*. 1996;335:384-91.
224. El-Sadr WM, Burman WJ, Grant LB, Matts JP, Hafner R, Crane L, et al. Discontinuation of prophylaxis for *Mycobacterium avium* complex disease in HIV-infected patients who have a response to antiretroviral therapy. *N Engl J Med*. 2000;342:1085-92.
225. Aberg JA, Yajko DM, Jacobson MA. Eradication of AIDS-related disseminated *Mycobacterium avium* complex infection after 12 months of antimycobacterial therapy combined with highly active antiretroviral therapy. *J Infect Dis*. 1998;178:1446-9.
226. Martínez E, Miró JM, González J, Mallolas J, Gatell JM. Withdrawal of *Mycobacterium avium* complex suppressive therapy in HIV-1-infected patients on highly active antiretroviral therapy. *Aids*. 1999;13:147-8.
227. Shafran SD, Gill MJ, Lalonde RG, Walmsley S-L, Toma E, Conway B, et al. Successful Discontinuation of MAC Therapy Following Effective HAART. Abstract 547. En: 8th Conference on Retroviruses and Opportunistic Infections; 2001; Chicago, IL; 2001.
228. Feikin DR, Feldman C, Schuchat A, Janoff EN. Global strategies to prevent bacterial pneumonia in adults with HIV-disease. *Lancet Infect Dis*. 2004;4:445-55.

229. Grau I, Pallares R, Tubau F, Schulze MH, Llopis F, Podzamcazer D, et al. Epidemiological changes in bacteremic pneumococcal disease in patients with human immunodeficiency virus in the era of highly active antiretroviral therapy. *Arch Intern Med.* 2005;165:1533-40.
230. French N, Nakiyingi J, Carpenter IM, Lugada E, Watera C, Moi K, et al. 23-valent pneumococcal polysaccharide vaccine in HIV-1-infected Ugandan adults: Double blind, randomised and placebo controlled trial. *Lancet.* 2000;355:2106-11.
231. Watera C, Nakiyingi J, Miro G, Nuwonge R, Whitworth JA, Gilks CF, et al. 23-valent pneumococcal polysaccharide vaccine in HIV-infected Ugandan adults: 6-year follow-up of a clinical trial cohort. *AIDS.* 2004;18:1210-3.
232. Peñaranda M, Falco V, Payeras A, Jordano Q, Curran A, Pareja A, et al. Effectiveness of polysaccharide pneumococcal vaccine in HIV-infected patients: A case-control study. *Clin Infect Dis.* 2007;45:e82-7.
233. Klugman KP, Madhi SA, Huebner RE, Kohberger R, Mbelle N, Pierce N, et al. A trial of a 9-valent pneumococcal conjugate vaccine in children with and those without HIV infection. *N Engl J Med.* 2003;349:1241-8.
234. Flannery B, Heffernan R, Harrison LH, Ray SM, Reingold AL, Hadler J, et al. Changes in invasive pneumococcal disease among HIV-infected adults living in the era of childhood pneumococcal immunisation. *Ann Intern Med.* 2006;144:1-9.
235. Gordin FM, Roediger MP, Girard PM, Lundgren JD, Miró JM, Palfreeman A, et al. Bacterial Pneumonia in HIV-Infected Persons: Increased Risk Associated with Cigarette Smoking and Treatment Interruption—Results from the SMART Study. *Am J Respir Crit Care Med.* 2008 [en prensa].
236. Gathiram V, Jackson TF. A longitudinal study of asymptomatic carriers of pathogenic zymodemes of *Entamoeba histolytica*. *S Afr Med J.* 1987;72:669-72.
237. McAuley JB, Herwaldt BL, Stokes SL, Becher JA, Roberts JM, Michelson MK, et al. Diloxanide furate for treating asymptomatic *Entamoeba histolytica* cyst passers: 14 years' experience in the United States. *Clin Infect Dis.* 1992;15:464-8.
238. Fung HB, Doan TL. Tinidazole: a nitroimidazole antiprotozoal agent. *Clin Ther.* 2005;27:1859-84.
239. Pehrson P, Bengtsson E. Treatment of non-invasive amoebiasis—a comparison between tinidazole and metronidazole. *Ann Trop Med Parasitol.* 1984;78:505-8.
240. Pehrson PO, Bengtsson E. A long-term follow up study of amoebiasis treated with metronidazole. *Scand J Infect Dis.* 1984;16:195-8.
241. Anonimo. Drug for Parasitic Infections. The Medical Letter. 2007;5 Suppl: 1-15.
242. Brentlinger PE, Behrens CB, Micek MA. Challenges in the concurrent management of malaria and HIV in pregnancy in sub-Saharan Africa. *Lancet Infect Dis.* 2006;6:100-11.
243. Idemiyor V. Human immunodeficiency virus (HIV) and malaria interaction in sub-Saharan Africa: the collision of two Titans. *HIV Clin Trials.* 2007;8:246-53.
244. Seed CR, Kitchen A, Davis TM. The current status and potential role of laboratory testing to prevent transfusion-transmitted malaria. *Transfus Med Rev.* 2005;19:229-40.
245. Arquin PM, Kozarsky PE, Reed C. Centers for Diseases Control. Health Information for International Travel 2008: US Department of Health and Human Services, Public Health Service; 2007.
246. World Health Organization. International Travel and Health; 2007.
247. Khoo S, Back D, Winstanley P. The potential for interactions between antimalarial and antiretroviral drugs. *Aids.* 2005;19:995-1005.
248. Tuset M, Miró JM, Codina C, Ribas J. Interacciones en VIH [consultado 30/07/2008]. Disponible en: <http://www.interaccionesshiv.com>
249. Schippers EF, Hugen PW, den Hartigh J, Burger DM, Hoetelmans RM, Visser LG, et al. No drug-drug interaction between nelfinavir or indinavir and mefloquine in HIV-1-infected patients. *Aids.* 2000;14:2794-5.
250. German P, S Parikh S, Lawrence J, Lindegardh N, Rosenthal P, Havlir D, et al. Drug Interaction between Antimalarial Drugs and Lopinavir/Ritonavir. Oral abstract 132. Paper presented at: 15th Conference on Retrovirus and Opportunistic Infections. Boston, February 3-6, 2008.
251. Cahn P, Badaró R, Freilij H. Other parasitic infections. En: Crowe S, Hoy J, Mills J, editors. Management of the HIV-infected patient. Chapter 31. 2nd ed. UK: Martin Dunitz Editors; 2002. p. 513-26.
252. Ferreira MS, Nishioka Sde A, Silvestre MT, Borges AS, Nunes-Araujo FR, Rocha A. Reactivation of Chagas' disease in patients with AIDS: report of three new cases and review of the literature. *Clin Infect Dis.* 1997;25:1397-400.
253. Silva N, O'Bryan L, Medeiros E, Holand H, Suleiman J, de Mendonca JS, et al. Trypanosoma cruzi meningoencephalitis in HIV-infected patients. *J Acquir Immune Defic Syndr Hum Retrovirol.* 1999;20:342-9.
254. Sartori AM, Lopes MH, Benvenuti LA, Caramelli B, di Pietro A, Nunes EV, et al. Reactivation of Chagas' disease in a human immunodeficiency virus-infected patient leading to severe heart disease with a late positive direct microscopic examination of the blood. *Am J Trop Med Hyg.* 1998;59:784-6.
255. Sartori AM, Shikanai-Yasuda MA, Amato Neto V, Lopes MH. Follow-up of 18 patients with human immunodeficiency virus infection and chronic Chagas' disease, with reactivation of Chagas' disease causing cardiac disease in three patients. *Clin Infect Dis.* 1998;26:177-9.
256. Yoo TW, Mlikotic A, Cornford ME, Beck CK. Concurrent cerebral american trypanosomiasis and toxoplasmosis in a patient with AIDS. *Clin Infect Dis.* 2004;39:e30-34.
257. Nishioka Sde A, Ferreira MS, Rocha A, Burgarelli MK, Silva AM, Duarte MI, et al. Reactivation of Chagas' disease successfully treated with benznidazole in a patient with acquired immunodeficiency syndrome. *Mem Inst Oswaldo Cruz.* 1993;88:493-6.
258. Solari A, Saavedra H, Sepulveda C, Oddo D, Acuna G, Labarca J, et al. Successful treatment of *Trypanosoma cruzi* encephalitis in a patient with hemophilia and AIDS. *Clin Infect Dis.* 1993;16:255-9.
259. Blum J, Nkunku S, Burri C. Clinical description of encephalopathic syndromes and risk factors for their occurrence and outcome during melarsoprol treatment of human African trypanosomiasis. *Trop Med Int Health.* 2001;6:390-400.
260. Louis JP, Moulija-Pelat JP, Jannin J, Asonganyi T, Hengy C, Trebucq A, et al. Absence of epidemiological inter-relations between HIV infection and African human trypanosomiasis in central Africa. *Trop Med Parasitol.* 1991;42:155.
261. World Health Organization: Control and surveillance of African trypanosomiasis. WHO Technical Report Series No 881. Geneva; 1998.
262. Fairlamb AH. Chemotherapy of human African trypanosomiasis: current and future prospects. *Trends in Parasitology.* 2003;19:488-94.
263. Pepin J, Khonde N, Maiso F, Doua F, Jaffar S, Ngampo S, et al. Short-course eflornithine in Gambian trypanosomiasis: a multicentre randomized controlled trial. *Bull World Health Organ.* 2000;78:1284-95.
264. Burri C, Nkunku S, Merolle A, Smith T, Blum J, Brun R. Efficacy of new, concise schedule for melarsoprol in treatment of sleeping sickness caused by *Trypanosoma brucei gambiense*: a randomised trial. *Lancet.* 2000;355:1419-25.
265. Pepin J, Mpia B. Randomized controlled trial of three regimens of melarsoprol in the treatment of *Trypanosoma brucei gambiense* trypanosomiasis. *Trans R Soc Trop Med Hyg.* 2006; 100:437-41.
266. Schmid C, Nkunku S, Merolle A, Voumatsou P, Burri C. Efficacy of 10-day melarsoprol schedule 2 years after treatment for late-stage gambiense sleeping sickness. *Lancet.* 2004;364:789-90.
267. Pape JW, Verdier R-I, Bony M, Bony J, Johnson WD. Cyclospora infection in adults infected with HIV: clinical manifestations, treatment, and prophylaxis. *Ann Intern Med.* 1994;121:654-7.
268. Keiser PB, Nutman TB. *Strongyloides stercoralis* in the immunocompromised population. *Clin Microbiol Rev.* 2004;17:208-17.
269. Viney ME, Brown M, Omoding NE, Bailey JW, Gardner MP, Roberts E, et al. Why does HIV infection not lead to disseminated strongyloidiasis? *J Infect Dis.* 2004;190:2175-80.
270. Page WA, Dempsey K, McCarthy JS. Utility of serological follow-up of chronic strongyloidiasis after anthelmintic chemotherapy. *Trans R Soc Trop Med Hyg.* 2006;100:1056.
271. Karunajeewa H, Kelly H, Leslie D, Leydon J, Saykao P, Biggs B-A. Parasite-specific igg response and peripheral blood eosinophil count following albendazole treatment for presumed chronic strongyloidiasis. *J Travel Med.* 2006;13:84-91.
272. Singthong S, Intapan PM, Wongsaroji T, Maleewong W. Randomized comparative trial of two high-dose albendazole regimens for uncomplicated human strongyloidiasis. *Southeast Asian J Trop Med Public Health.* 2006; 37 Suppl 3:32-4.
273. Zaha O, Hirata T, Uchima N, Kinjo F, Saito A. Comparison of anthelmintic effects of two doses of ivermectin on intestinal strongyloidiasis in patients negative or positive for anti-HTLV-1 antibody. *J Infect Chemother.* 2004;10:348-51.
274. Marti H, Haji HJ, Savioli L, Chwaya HM, Mgeni AF, Ameir JS, et al. A comparative trial of a single-dose ivermectin versus three days of albendazole for treatment of *Strongyloides stercoralis* and other soil-transmitted helminth infections in children. *Am J Trop Med Hyg.* 1996;55:477-81.
275. Toma H, Sato Y, Shiroma Y, Kobayashi J, Shimabukuro I, Takara M. Comparative studies on the efficacy of three anthelmintics on treatment of human strongyloidiasis in Okinawa, Japan. *Southeast Asian J Trop Med Public Health.* 2000;31:147-51.
276. King CH, Mahmoud AA. Drugs five years later: praziquantel. *Ann Intern Med.* 1989;110:290-6.
277. Grandiere-Pérez L, Ansart S, Paris L, Faussart A, Jauraguiberry S, Grivois JP, et al. Efficacy of praziquantel during the incubation and invasive phase of *Schistosoma haematobium* schistosomiasis in 18 travelers. *Am J Trop Med Hyg.* 2006;74:814-8.

278. Serpa JA, Moran A, Goodman JC, Giordano TP, White AC, Jr. Neurocysticercosis in the HIV era: a case report and review of the literature. *Am J Trop Med Hyg.* 2007;77:113-7.
279. Prasad S, MacGregor RR, Tebas P, Rodriguez LB, Bustos JA, White AC, Jr. Management of potential neurocysticercosis in patients with HIV infection. *Clin Infect Dis.* 2006;42:e30-34.
280. Singhi P, Ray M, Singhi S, Khandelwal N. Clinical spectrum of 500 children with neurocysticercosis and response to albendazole therapy. *J Child Neurol.* 2000;15:207-13.
281. Kalra V, Dua T, Kumar V. Efficacy of albendazole and short-course dexamethasone treatment in children with 1 or 2 ring-enhancing lesions of neurocysticercosis: A randomized controlled trial. *J Pediatrics.* 2003;143:111-4.
282. Proano JV, Madrazo I, Avelar F, López-Felix B, Díaz G, Grijalva I. Medical treatment for neurocysticercosis characterized by giant subarachnoid cysts. *N Engl J Med.* 2001;345:879-85.
283. Garcia HH, Pretell EJ, Gilman RH, Martínez SM, Moulton LH, Del Brutto OH, et al. A trial of antiparasitic treatment to reduce the rate of seizures due to cerebral cysticercosis. *N Engl J Med.* 2004;350:249-58.
284. López-Gómez M, Castro N, Jung H, Sotelo J, Corona T. Optimization of the single-day praziquantel therapy for neurocysticercosis. *Neurology.* 2001;57:1929-30.
285. Del Brutto OH, Roos KL, Coffey CS, García HH. Meta-Analysis: cysticidal drugs for neurocysticercosis: albendazole and praziquantel. *Ann Intern Med.* 2006;145:43-51.
286. Góngora-Rivera F, Soto-Hernández JL, González Esquivel D, Cook HJ, Márquez-Caraveo C, Hernández Dávila R, Santos-Zambrano J. Albendazole trial at 15 or 30 mg/kg/day for subarachnoid and intraventricular cysticercosis. *Neurology.* 2006;66:436-8.
287. Nutman TB, Miller KD, Mulligan M, Reinhardt GN, Currie BJ, Steel C, Ottesen EA. Diethylcarbamazine prophylaxis for human loiasis. Results of a double-blind study. *N Engl J Med.* 1988;319:752-6.